

# Acetaminophen Dosing (For Example, Tylenol®)

For infants and children OLDER than 2 months old.  
Give every 4-6 hours as needed for fever or pain. Do not give any other medications that contain acetaminophen

NOTE: Certain concentrations (mg/mL) of acetaminophen are being phased out.  
Please be sure to check the concentration on the bottle before dosing.

WEIGHT (Pounds)	AGE	CHILDREN'S LIQUID (160 mg / 5 mL)	CHILDREN'S TABS (160 mg Tabs)
6-11 lbs	0-3 months	1.25 mL (1/4 tsp)	—
12-17 lbs	4-11 months	2.5 mL (1/2 tsp)	—
18-23 lbs	12-23 months	3.75 mL (3/4 tsp)	—
24-35 lbs	2-3 years	5 mL (1 tsp)	1 Tab
36-47 lbs	4-5 years	7.5 mL (1 1/2 tsp)	1.5 Tabs
48-59 lbs	6-8 years	10 mL (2 tsp)	2 Tabs
60-71 lbs	9-10 years	12.5 mL (2 1/2 tsp)	2.5 Tabs
72-95 lbs	11 years	15 mL (3 tsp)	3 Tabs



# Ibuprofen Dosing (For Example, Motrin® and Advil®)

For infants and children OLDER than 6 months of age.  
Give every 6-8 hours as needed for fever or pain.  
DO NOT GIVE MORE THAN 4 DOSES IN 24 HOURS.  
Do NOT use with any other medicine containing ibuprofen.

WEIGHT (Pounds)	AGE	CHILDREN'S LIQUID (100 mg / 5 mL)	CHILDREN'S TABS (50 mg Tabs)
0-11 lbs	0-5 months	—	—
12-17 lbs	6-11 months	2.5 mL	—
18-23 lbs	12-23 months	4 mL	—
24-35 lbs	2-3 years	5 mL (1 tsp)	2 Tab
36-47 lbs	4-5 years	7.5 mL (1 1/2 tsp)	3 Tabs
48-59 lbs	6-8 years	10 mL (2 tsp)	4 Tabs
60-71 lbs	9-10 years	12.5 mL (2 1/2 tsp)	5 Tabs
72-95 lbs	11 years	15 mL (3 tsp)	6 Tabs
96 lbs or more	12 years or older	20 mL (4 tsp)	8 Tabs